



# MEDICINAL HERBS AFFECTING GONADOTROPIN HORMONES IN WOMEN:AN UPDATED SYSTEMATIC REVIEW

FARINAZ FARAHBOD<sup>1</sup>, SAEID HEIDARI SOURESHJANI<sup>2\*</sup>

<sup>1</sup>Assistant Professor of Obstetrics & Gynecology, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup> MSc, research assistant, Deputy of Research and Technology, Shahrekord University of Medical Sciences, Shahrekord, Iran.

## ABSTRACT

Plants and their derivatives can exacerbate or relieve diseases such as infertility, puberty disorders, and menopause. The present - study examines the role and mechanism of medicinal plants or their products in reducing or increasing the levels of GnRH, FSH and LH in females. The key word "GnRH" or "FSH" or "Luteinizing Hormone" was searched in conjunction with the words Medicinal plant, Herb \* and Phyto \* in the ISI and PubMed databases (using Endnote software). Then articles related to herbs and products affecting the hormones of GnRH, FSH and LH, were included in the study, following inclusion and exclusion criteria. Plants and plant derivatives that affect fertility disorders and mainly increase GnRH, include *Vitex agnus-castus*, *Thuja occidentalis* L., *Cimicifuga racemosa*, *Yucca schidigera*, isoflavones and some Chinese herbal compounds. In some cases, plants such as *Emilia coccinea* decrease the fertility of the fetus by lowering the FSH and LH hormones. Also, some herbs and plant compounds are used for disorders and discontinuation of symptoms. These include isoflavones, *Lepidium meyenii*, *Angelica gigas* and Chinese herbal compounds. In some cases, *Radix Ginseng* and isoflavones also cause premature aging and puberty, which this issue should be taken into account for nutritional purposes. Plants and plant compounds, can cause hormonal changes, especially in female GnRH hormones mainly due to presence of having phytoestrogens and by inducing their estrogenic properties.

**KEYWORDS:** GnRH, female, medicinal herbs, sexual disorders

## INTRODUCTION

The medicinal plants are used due to the least side effects and lower prices for the treatment of psychological and physiological diseases<sup>1-8</sup>. Today, despite the existence of multiple treatments to promote different dimensions of health, chemical and herbal treatments are still of particular importance in treating various disorders and diseases<sup>9-12</sup>. The prevalence of infertility among young women in various countries is increasing<sup>13-16</sup>. Chemical treatments and infertility treatments impose a lot of expense on patients as well as trigger many psychological and emotional stress to them and the complications of these therapies may have an adverse effect on the patients' health<sup>17,18</sup>. The balance of hormones in the body plays a major role in successful pregnancy, or prevents the onset of the reproductive system The

gonadotropin releasing hormone (GnRH) is one of the hormones that plays an important role in women's reproduction, and the sex hormones (FH) of the Follicle-Stimulating Hormone and (LH) Luteinizing Hormone release from the pituitary gland in response to GnRH<sup>19</sup>. These hormones stimulate ovulation and normalize oocyte growth in women<sup>20</sup>. On the other hand, hormonal disorders also cause symptoms such as hot flashes and other complications in menopause. LH hormone produces less estrogen. Although, the level of FSH still rises after the menopause, eventually the level decreases. Therefore, hormonal balance in this age of menopause is very important<sup>21,22</sup>. Modulation of these hormones can help in the treatment of infertility and hormonal disorders in women. It also helps in Therapeutic methods including assisted reproductive technology (ART) and treatment of disorders such as polycystic ovary syndrome<sup>23</sup>, ovarian hyperstimulation syndrome<sup>24</sup> and

hypogonadotropic hypogonadism<sup>20</sup>. Studies have shown that medicinal herbs can be effective treatments for various diseases, including sexual disorders<sup>25</sup>. Therefore, due to the increasing use of medicinal plants and the importance of hormonal balance in the treatment of infertility, a systematic study has been conducted to investigate the role and mechanism of medicinal plants or their products in reducing or increasing the levels of GnRH, FSH and LH in female.

## METHODS

This systematic study was performed by inserting the keywords in the Endnote software. The key terms were searched in The GnRH, or "FSH" or "Luteinizing Hormone" in combination with the words Medicinal plant, Herb \* and Phyto \* in the ISI and PubMed Databases. Having identified the available articles and documents, the abstracts of the articles, as well as information from other sources by the researchers, herbs and products that influenced the hormones of GnRH, FSH and LH were selected. Also, merely clinical trials and laboratory studies between 2010 and 2017 were studied and investigated. Articles which did not have full text was not available and those that were not related to the purpose of the study were excluded from the study. Figure 1 illustrates exclusion and inclusion criteria and, studies that meet the inclusion criteria

## RESULTS

Plants, herbal derivatives, and herbal formulations shown to be able to induce endocrine function and influence sex hormones in different way.in this study, we reviewed this issue systematically.

### *Medicinal Plants*

#### *Vitex agnus-castus*

This plant, which has antioxidant properties and is a phytoestrogen, has been investigated on hypothalamic-pituitary-gonadal axis in experimental mice. The study showed that the gavage of laboratory mice with the extract of this plant can increase the level of LH and FSH hormones. Thus, the Vitex extract enhances the fertility power of old female mice<sup>27</sup>.

#### *Angelica gigas*

This is one of the herbal medicines used in butter. The root powder of this plant regulates the levels of ovarian hormones and possesses anti-osteoporosis

properties. The results of the study indicated that the levels of FSH hormone decreased in the rats that were administered Korean angelica powder, but the serum levels of LH in laboratory rats increased<sup>28</sup>.

#### *Renshen (Radix Ginseng)*

The effect of estrogenic Renshen on reproductive tissues in immature mice was evaluated. Extract of this plant stimulates estrogen biosynthesis and has a negative effect on the level of LH and FSH hormones<sup>29</sup>.

#### *Emilia coccinea*

This plant is used in Africa as a contraceptive. Doses of 1000 and 100 mg / kg / day on mice that were not pregnant were evaluated to determine sexual function and hormonal changes. In this study, high-dose extract reduced the level of FSH and LH in serum without significant changes in estrogen and progesterone levels and ultimately prevented ovulation<sup>30</sup>.

#### *Thuja occidentalis L. (Cupressaceae)*

T. occidentalis are used in various diseases of women, to treat polycystic ovary syndrome (PCOS) in laboratory rats. In this study, the oil of this plant with  $\alpha$ -thujone after 21 days of administration has been shown to reduce LH and may be used as an effective treatment (without induction of osteoporosis) in the treatment of PCOS<sup>31</sup>.

#### *Cimicifuga racemosa*

It contains phytoestrogens together with clomiphene in women with PCOS lowers LH levels, but does not alter FSH levels. Thus, adding the extract of this plant to clomiphene increases the chances of fertility in women with PCOS by inducing hormonal changes such as increased levels of estradiol and progesterone and increasing the thickness of the endometrium and producing more follicles<sup>32</sup>.

#### *Yucca schidigera*

The addition of this herbal supplement to sheep nutrition increases ovarian response to FSH and, improves folliculogenesis by inducing other cellular changes in the ovaries and thus increases the likelihood of fertility<sup>33</sup>.

#### *Lepidium meyenii (Maca)*

Maca is used in disabilities and hormonal disorders. Hydroalcoholic extract of this plant in ovariectomized mice resulted in elevated FSH and estradiol levels even at low doses. Therefore, the

plant can be a natural therapeutic choice for postmenopausal women<sup>34</sup>.

### Plant derivative

#### Isoflavones

Because of these phytoestrogenic properties, Isoflavones in soybean can make a variety of changes to the hormonal system. A study on postmenopausal women showed that supplementation with 70 mg per day of isoflavone after 12 weeks reduced LH levels in women, but did not change levels of FSH, estrone and estradiol<sup>35</sup>. Exposure of newborn rats to soy isoflavones have increased FSH levels and early exposure to these estrogenic compounds can impair the maturity of these animals<sup>36</sup>. Genistein is considered as an isoflavone and is a phytoestrogens, which was studied in Bhattarai et al. Its effects on gonadotrophin releasing hormone neurons in young female mice was evaluated. The results showed that this plant phytoestrogen induces GABA and kisspeptin-mediated transmissibility to GnRH neurons with its stimulatory activity on GnRH neurons and thus can increase fertility in laboratory rats.<sup>37</sup> Genistein significantly affects the expression of the analyzed estrogen-response genes and reduces the LH and FSH hormones in the rats, which is probably due to the estrogenic properties of the substance as well as the effect on the release of gonadotrophins<sup>38</sup>. Genistein has dose-dependent effects that increase estradiol in the mean to high dose and reduce both the LH and FSH hormones and thus alleviate the symptoms of abnormal ovarian histomorphology. Hence, Genistein is effective in treating premature ovarian failure and subsequent infertility treatment<sup>39</sup>. Genistein increases the number of pituitary gonadotropic cells in addition to the changes in the hormones in a way that increase the number of FSH cells (by 21%) and LH cells (by 20%) per mm<sup>2</sup><sup>40</sup>. In gilts, exposure to Genistein also leads to changes in endocrine glands, including lowering LH levels after artificial insemination<sup>41</sup>. Also, exposure to this plant phytoestrogen during infancy (rats) can disrupt the normal development of the neuroendocrine system and increase serum FSH levels at higher concentrations of the drug but lower LH levels<sup>42</sup>. *Isoflavone daidzein* is one of the other phytoestrogens whose effect on the function of the cells of basic porcine ovarian granulosa and the response to FSH was investigated. The results showed that FSH stimulates the proliferation of ovarian cells and modifies the daidzein of this hormone and thus has a direct effect on ovaries and women's fertility<sup>43</sup>.

### Herbal compounds and formulation

#### Chinese herbal medicine formula

Combined drugs used in the treatment of infertility may be used with or without acupuncture in traditional Chinese medicine. For example, in a study by Hullender et al.<sup>43</sup>, a combination of Tao Ren 20% (*Persicae* spp. Semen), Hong Hua 20% (*Carthami* spp. Flos), Chuan Xiong 10% (*Chuanxiong* rhizoma), Shu di Huang 20% (*Rehmannia preparata* radix), Dang Gui 20% (*Angelica sinensis* radix) and Bai Shao 10% (*Paeoniae alba* radix) were used along with acupuncture to treat a patient with diminished ovarian reserve (DOR). The patient's FSH level changed from 14.5 mIU / mL to 8.7 mIU / mL after treatment, and after 3 cycles and more with clomid, the level increased to 16.8 mIU / mL<sup>44</sup>. Lirukang Granule was used with psychotherapy in patients with cyclomastopathy and menoxenia. This combination includes plants such as *Radix Curcumae*, *pseudobarbus Cremstrae seu Pleiones*, *Colla Cornus Cervi* and *Concha Ostreae* at 12 grams per sac. After treatment with this plant, FSH and other sex hormones such as estradiol, progesterone and prolactin increased<sup>45</sup>. In a similar study that investigated the effects of Xiaoyao Pill and Kunbao Pill twice a day on peri-menopausal syndrome and hyperlipidemia, these two plant compounds reduced the level of LH and FSH, thus maintaining hormonal balance in individuals Studied<sup>46</sup>. Su et al. Used *Radix Astragali*, *Radix Angelicae Sinensis*, and *Folium Epimedii* to reduce peri-monosuppression symptoms in rats. After administration of these compounds, estradiol increased in circulation, but the level of FSH decreased. However, changes in LH levels were not observed, and thus this Chinese herbal composition can be used to modulate the sex hormones in the peri-monoplasal syndrome<sup>47</sup>. Shouwu jiangqi is one of the Chinese drug compounds whose effect of extract was studied in Wang et al. on PCOS mice. Extract of *shouwu jiangqi* in high doses with dimethylbiguanide increases the level of FSH but does not change the amount of LH. By altering the plant's composition, this plant Reduces ovarian dysfunction in mice<sup>48</sup>. Acetylshikonin is a plant compound that is a derivative of Zicao (Chinese herbal remedy). Acetylshikonin in laboratory rats reduces LH and FSH levels by inhibiting the secretion or synthesis of these hormones. This is due to its effect on secretion of the exocytosis process of gonadotropic hormone, which leads to infertility in rats<sup>49</sup>. *Yiru Tiaojing* Granule is a Chinese herbal compound consisting of *Rhizoma Curculiginis*

27.2%, *Radix Paeoniae alba* 27.2%, *Radix Morinda officinalis* 22.7%, and *Radix Glycyrrhizae* 22.7%. In the study of the effect of reduction of prolactin *Yiru Tiaojing* Granule, has been shown to affect brain receptors and prevent hyperprolactinemia in rat models. It was also found to increase the levels of LH, estradiol, and progesterone hormones<sup>50</sup>. In a study by Wu et al., a plant compound composed of *Radix Pseudostellariae* 30 g, *Rhizoma Polygonati Odorati* 15 g, *Radices Paeoniae Alba* 12 g, *Eucommia ulmoides* 12 g, *Cornus officinalis* 12 g, *Ligustrum lucidum* 15 g, *Fructus Mume* 12 g, *Semen cuscuteae* 15 g, *Radix Pueraria* 10 g and *Radix Trichosanthis* 12 g, *Fructus Lycii* 15 g and *periostracum cicada* 6 g, *Radix Clematidis* 15 g, *Herba Siegesbeckiae* 15 g, *Herba Sarcandrae* 12 g and *Herba Lycopodti* 15 g w *Corydalis Tuber* 12 g, *Finger Citron Fructus Citri Sarcodactylis* 12 g, *Rhodiola dodder* 15 g and *Fructus Mori* 15 g.) To Nourishing yin, supplementing qi, and activating blood were used. Each of These compounds used to address the problem of patients with Primary Sjogren's Syndrome, led to the regulation of reproductive endocrine-immune network. It decreased levels of LH and FSH and regulated other hormones and factors that were elevated abnormally<sup>51</sup>. *Bu Shen Tiao Chong* recipe was used to study the molecular pathway in retrieving diminished ovary reserve. This compound consist of medicinal Indian mulberry root (10 g), *Szechuan lovage rhizome* (6 g), Chinese angelica (10 g), mayflower Solomon's seal rhizome (15 g), desertliving *cistanche* (10 g), prepared rehmannia root (15 g), South Dodder seed (15 g), Chinese magnolia vine fruit (6 g), and epimedium herb and fluorite (15 g). It was found that the *Bu Shen Tiao Chong* recipe produces hormones through the equilibrium that it may have a beneficial role in oocyte maturation and repair of the diminished ovary reserve. The plant extract plays this role through the restoring of the depleted brain-derived neurotrophic factor and up regulating the FSH, E2, and Inhibin B hormones<sup>52</sup>. *Radix Astragali*, *Radix Angelicae Sinensis*, and *Folium Epi- medii* are compounds of the medicinal plants used to treat sexual problems. The combination was evaluated for ovariectomized rats and natural aging female rats to determine their safety and efficacy. In this study, it was found that low levels of FSH and LH and other sex hormones were regulated and could be used to treat perimenopausal symptoms. But long-term use of estrogenic effects should be considered<sup>53</sup>. *QiBaoMeiRan* formula is a combination of *Polygoni Multiflori Radix*, *Angelicae Sinensis Radix*, *Achyranthis Bidentatae*

*Radix*, *Semen Cuscuteae*, *Fructus Lycii*, *Poria*, and *Fructus Psoraleae*). Due to its phytoestrogenic properties and elevated levels of estradiol decreases the level of FSH and LH in immature rats<sup>54</sup>. The study of Zeng et al.<sup>55</sup> showed that the use of Chinese herbal combination named Yin, could be effective in decreasing the expression of hypothalamic mTOR mRNA and p-mTOR protein and lowering serum levels of LH and estradiol, thereby delaying maturation of mice. A vivo model of ovariectomized plant compound called Qing'E was studied. This combined powder contains *Eucommniae Cortex* (480 g, beaked with salt), *Psoraleae Fructus* (240 g, beaked with salt), *Juglandis Semen* (150 g), and *Garlic Rhizoma* (120 g, steamed and dried) which, like other compounds, this plant caused elevated levels of estradiol and subsequently LH due to phytoestrogens<sup>56</sup>. Consumption of Chinese Herbs in the following doses of 20g of *Radix Astragali*, 35g of *Radix Rehmanniae Preparata*, 30g of *Cuscuta chinensis Lam*, 35g of *Fructus Ligustri Lucidi*, 30g of *Fructus Psoraleae*, 20g of *Radix Salviae Miltiorrhizae* and 20g of *Rubus idaeus Linn* in combination with other therapeutic methods like Acupuncture can increase the efficiency of these plants. It causes the decrease in the level of FSH and LH in mice with PCOS<sup>57</sup>. Medicinal plants can affect the regulation and modulation of sex hormones in women through binding affinity for estrogen receptors, directly and indirectly by induction of phytoestrogenic properties. Medicinal herbs or their derivatives have several mechanisms for modulating the gonadotropin hormones, as summarized in Fig. 2<sup>33,52,55</sup>. Figure 1: Flowchart related to the procedure of study of studies. Figure 2: Mechanisms of Effect of Medicinal Herbs and Their Compounds on GnRH Hormonal modulation. Also, medicinal plants can increase estradiol serum levels due to their high estrogenic properties in high doses and in prolonged usage. This is due to the presence of plant phytoestrogens that, with elevated levels of estradiol, result in negative feedback in the hypothalamus, leading to a decrease in GnRH production<sup>54,58</sup>.

## DISCUSSION

The balance of sexual hormones in humans is very important. To produce a successful fertility in the first half of the menstrual cycle, estrogen levels rise and cause the growth and increase of uterine cells. In response to the FSH hormone, an ovum begins to grow inside one of the ovaries. On day of 14 of a 28-day cycle, an ovary is released in response to the

LH hormone. And the increase in progesterone levels occurs in the second half of the menstrual cycle, which results in an increase in the thickness of the lower limb<sup>32,59</sup>. Therefore, the time taken to use herbal drugs in this interval (the first half of the cycle until the 14th day of a 28-day cycle) is more important for increasing the quantity and quality of oocytes. What should be considered is that in postmenopausal women, levels of LH and FSH levels increase<sup>60,61</sup>. And the results of the studies may be influenced by confounding factors such as age, weight and lifestyle, and other underlying conditions (such as those that reduce the function of the ovaries)<sup>28</sup>. Also, poor responder women report weak response to gonadotropins<sup>62</sup>, which should also be considered in plant treatments. Also, soybean isoflavones in animals during puberty or pre-puberty period have adverse effects and disrupt the hormonal balance in these animals. It increases FSH but lowers LH and estradiol levels. These changes, along with other problems, such as down regulation, can cause genes that play a key role in the synthesis of steroid hormones and delay in puberty, which can disrupt the normal functioning of sexual cells<sup>63,64</sup>. In infancy, due to exposure to phytoestrogens such as Genistein, a disorder occurs in the development of the neuroendocrine system that should be addressed in intake of foods such as soybeans or other herbal supplements<sup>42</sup>. It should

be noted that consuming some plants during fertility can have adverse effects on the fetus or mother and may lead to abortion and teratogenic side effects. On the other hand, some herbs or herbal compounds may lower the LH and FSH hormones and cause other changes in hormones and enzymes<sup>65</sup>; as an example, Plant-derived nutritional supplements can, in addition to the formation of toxicity, reduce the level of LH and FSH hormones and disrupt the hormonal balance<sup>66</sup>. Hence, recognizing plant chemical properties or consulting with a doctor or specialist can be an important step in the treatment of infertility. Plants and plant compounds can cause hormonal changes, especially in female GnRH hormones, mainly due to phytoestrogens and by their estrogenic properties induction. These hormonal changes may be beneficial or harmful based on age, sex life, and type of sexual dysfunction. Therefore, the necessary precautions in this regard should be considered. On the other hand, the dosage and duration of use of medicinal herbs are also considered as two important factors in the effectiveness of natural therapies.

## CONFLICT OF INTEREST

Conflict of interest declared none.

## REFERENCES

1. Shabanian S, Khalili S, Lorigooini Z, Malekpour A, Heidari-Soureshjani S: The effect of vaginal cream containing ginger in users of clotrimazole vaginal cream on vaginal candidiasis. *Journal of Advanced Pharmaceutical Technology & Research* 2017;8:80-84.
2. Yavangi M: Use of Iranian Medicinal Plants Effective on Male Fertility Indices. *Journal of Global Pharma Technology* 2017;8:36-43.
3. Yavangi M: A Systematic Review of Iranian Medicinal Plants Effective on Female Infertility. *Journal of Global Pharma Technology* 2017;8:44-49.
4. Nikfarjam M, Bahmani M, Heidari-Soureshjani S. Phytotherapy for anxiety in Iran: A review of the most important Anti-anxiety medicinal plants. *Journal of Chemical and Pharmaceutical Sciences* 2016; 9 (3). 1235-1241.
5. Nikfarjam M, Bahmani M, Heidari-Soureshjani S: Phytotherapy for depression: A review of the most important medicinal plants of flora of Iran effective on depression. *Journal of Chemical and Pharmaceutical Sciences* 2016; 9 (3). 1242-1247.
6. Heidari-Soureshjani S, Asadi-Samani M, Yang Q, Saeedi-Boroujeni A: Phytotherapy of nephrotoxicity-induced by cancer drugs: an updated review. *Journal of Nephropathology* 2017;6:254-263.
7. Asadi-Samani M, Raeisi R, Heidari-Soureshjani S, Asadi-Samani M: A review for discovering hepatoprotective herbal drugs with least side effects on kidney. *Journal of Nephrotherapeutics* 2017;6:38-48.
8. Solati K, Heidari-Soureshjani S, Luther T, Asadi-Samani M: Iranian medicinal plants effective on sexual disorders: A systematic review. *International journal of pharmaceutical sciences and research* 2017;8:2415-2420.
9. Nikfarjam M, Heidari-Soureshjani S, Khoshdel A, Asmand P, Ganji F: Comparison of spiritual well-being and social health among the students attending group and individual religious rites. *World Family*

- Medicine/Middle East journal of Family Medicine 2017;15:160-165.
10. Artimani T, Shabani S, Heidari-Soureshjani S, Asadi-Samani M, Luther T, Branch S: A review of Iranian medicinal plants with teratogenic and abortion-inducing side effects. *International journal of pharmaceutical sciences and research* 2017;8:2372-2377.
  11. Ardestani-Samani N, Rabiei M, Ghasemi-Pirbalooti M, Bayati A, Heidari-Soureshjani S: Comparative study of self-concept, physical selfconcept, and time perspective between the students with multiple sclerosis and healthy students in Shahrekord. *World Family Medicine/Middle East Journal of Family Medicine* 2017;15:80-84.
  12. Ardestani-Samani N, Rabiei M, Ghasemi-Pirbalooti M, Bayati A, Heidari-Soureshjani S: Study and comparison of psychological disorders in normal students and students with multiple sclerosis in Shahrekord. *World Family Medicine/Middle East Journal of Family Medicine* 2017;15:75-79.
  13. Kazemijalilseh H, Ramezani Tehrani F, Behboudi-Gandevani S, Hosseinpanah F, Khalili D, Azizi F. The Prevalence and Causes of Primary Infertility in Iran: A Population-Based Study. *Global journal of health science*. 2015;7(6):226-32.
  14. Akhondi MM, Kamali K, Ranjbar F, Shirzad M, Shafeghati S, Behjati Ardakani Z, et al. Prevalence of Primary Infertility in Iran in 2010. *Iranian Journal of Public Health*. 2013;42(12):1398-404.
  15. Bushnik T, Cook JL, Yuzpe AA, Tough S, Collins J. Estimating the prevalence of infertility in Canada. *Human Reproduction (Oxford, England)*. 2012;27(3):738-46.
  16. Gokler ME, Unsal A, Arslantas D. The Prevalence of Infertility and Loneliness among Women Aged 18-49 Years Who Are Living in Semi-Rural Areas in Western Turkey. *International Journal of Fertility & Sterility*. 2014;8(2):155-62.
  17. Katz P, Showstack J, Smith JF, Nachtigall RD, Millstein SG, Wing H, et al. Costs of infertility treatment: Results from an 18-month prospective cohort study. *Fertility and sterility*. 2011;95(3):915-21.
  18. Hasanpoor-Azghdy SB, Simbar M, Vedadhir A. The emotional-psychological consequences of infertility among infertile women seeking treatment: Results of a qualitative study. *Iranian Journal of Reproductive Medicine*. 2014;12(2):131-8.
  19. Schneider F, Tomek W, Grundker C. Gonadotropin-releasing hormone (GnRH) and its natural analogues: a review. *Theriogenology*. 2006;66(4):691-709.
  20. Raju GAR, Chavan R, Deenadayal M, Gunasheela D, Gutgutia R, Haripriya G, et al. Luteinizing hormone and follicle stimulating hormone synergy: A review of role in controlled ovarian hyper-stimulation. *Journal of Human Reproductive Sciences*. 2013;6(4):227-34.
  21. Santoro N, Randolph JF. Reproductive Hormones and the Menopause Transition. *Obstetrics and gynecology clinics of North America*. 2011;38(3):455-66.
  10. Butler L, Santoro N. The Reproductive Endocrinology of the Menopausal Transition. *Steroids*. 2011;76(7):627-35.
  22. Kumar P, Sait SF. Luteinizing hormone and its dilemma in ovulation induction. *Journal of Human Reproductive Sciences*. 2011;4(1):2-7.
  23. Cooper O, Geller JL, Melmed S. Ovarian hyperstimulation syndrome caused by an FSH-secreting pituitary adenoma. *Nature clinical practice Endocrinology & metabolism*. 2008;4(4):234-8.
  24. Yavangi M. Use of Iranian Medicinal Plants Effective on Male Fertility Indices. *Journal of Global Pharma Technology*. 2017;8(10):36-43.
  25. Yavangi M. A Systematic Review of Iranian Medicinal Plants Effective on Female Infertility. *Journal of Global Pharma Technology*. 2017;8(10):44-9.
  26. Ahangarpour A, Najimi SA, Farbood Y. Effects of Vitex agnus-castus fruit on sex hormones and antioxidant indices in a d-galactose-induced aging female mouse model. *Journal of the Chinese Medical Association : JCMSA*. 2016;79(11):589-96.
  27. Choi KO, Lee I, Paik SYR, Kim DE, Lim JD, Kang WS, et al. Ultrafine Angelica gigas Powder Normalizes Ovarian Hormone Levels and Has Antiosteoporosis Properties in Ovariectomized Rats: Particle Size Effect. *Journal of medicinal food*. 2012;15(10):863-72.
  28. Ding J, Xu Y, Ma XP, An JN, Yang XD, Liu ZQ, et al. Estrogenic effect of the extract of Renshen (Radix Ginseng) on reproductive tissues in immature mice. *Journal of*

- Traditional Chinese Medicine. 2015;35(4):460-7.
29. Elvis-Offiah UB, Bafor EE, Eze GI, Igbinumwen O, Viegelmann C, Edrada-Ebel R. In vivo investigation of female reproductive functions and parameters in nonpregnant mice models and mass spectrometric analysis of the methanol leaf extract of *Emilia Coccinea* (Sims) G Dons. *Physiol Rep*. 2016;4(23).
  30. Kupeli Akkol E, Ilhan M, Ayse Demirel M, Keles H, Tumen I, Suntar I. Thuja occidentalis L. and its active compound, alpha-thujone: Promising effects in the treatment of polycystic ovary syndrome without inducing osteoporosis. *Journal of ethnopharmacology*. 2015;168:25-30.
  31. Shahin AY, Mohammed SA. Adding the phytoestrogen *Cimicifugae Racemosae* to clomiphene induction cycles with timed intercourse in polycystic ovary syndrome improves cycle outcomes and pregnancy rates - a randomized trial. *Gynecological endocrinology : the official journal of the International Society of Gynecological Endocrinology*. 2014;30(7):505-10.
  32. Vlckova R, Sopkova D, Andrejckova Z, Valocky I, Kadasi A, Harrath AH, et al. Dietary supplementation of yucca (*Yucca schidigera*) affects ovine ovarian functions. *Theriogenology*. 2017;88:158-65.
  33. Zhang Y, Yu L, Jin W, Ao M. Effect of ethanolic extract of *Lepidium meyenii* Walp on serum hormone levels in ovariectomized rats. *Indian journal of pharmacology*. 2014;46(4):416-9.
  34. Kim J, Lee H, Lee O, Lee KH, Lee YB, Young KD, et al. Isoflavone supplementation influenced levels of triglyceride and luteinizing hormone in Korean postmenopausal women. *Archives of pharmacological research*. 2013;36(3):306-13.
  35. Muller DR, Soukup ST, Kurrat A, Liu X, Schmicke M, Xie MY, et al. Neonatal isoflavone exposure interferes with the reproductive system of female Wistar rats. *Toxicology letters*. 2016;262:39-48.
  36. Bhattarai JP, Abraham IM, Han SK. Genistein excitation of gonadotrophin-releasing hormone neurones in juvenile female mice. *Journal of neuroendocrinology*. 2013;25(5):497-505.
  37. Bliedtner A, Zierau O, Albrecht S, Liebhaber S, Vollmer G. Effects of genistein and estrogen receptor subtype-specific agonists in ArKO mice following different administration routes. *Molecular and cellular endocrinology*. 2010;314(1):41-52.
  38. Ding Q, Wang Y, Li N, Zhu K, Hu J, Wang S, et al. Bidirectional Estrogen-Like Effects of Genistein on Murine Experimental Autoimmune Ovarian Disease. *International journal of molecular sciences*. 2016;17(11).
  39. Medigovic I, Manojlovic-Stojanoski M, Trifunovic S, Ristic N, Milosevic V, Zikic D, et al. Effects of genistein on gonadotropic cells in immature female rats. *Acta Histochem*. 2012;114(3):270-5.
  40. Norrby M, Madsen M, Saravia F, Lundeheim N, Madej A. Genistein alters the release of oxytocin, prostaglandins, cortisol and LH during insemination in gilts. *Reproduction in domestic animals = Zuchthygiene*. 2011;46(2):316-24.
  41. Zin SRM, Omar SZ, Khan NLA, Musameh NI, Das S, Kassim NM. Effects of the phytoestrogen genistein on the development of the reproductive system of Sprague Dawley rats. *Clinics*. 2013;68(2):253-62.
  42. Sirotkin AV, Alexa R, Kadasi A, Stochmal'ova A, Morovic M, Laurincik J, et al. The isoflavone daidzein directly affects porcine ovarian cell functions and modifies the effect of follicle-stimulating hormone. *Journal of animal physiology and animal nutrition*. 2017;101(1):127-35.
  43. Hullender Rubin L, Marx BL. Diminished Ovarian Reserve, Clomid, and Traditional Chinese Medicine: A Case Study. *Medical acupuncture*. 2012;24(4):273-80.
  44. Bejaoui A, Chaabane H, Jemli M, Boulila A, Boussaid M. Essential oil composition and antibacterial activity of *Origanum vulgare* subsp. *glandulosum* Desf. at different phenological stages. *Journal of medicinal food*. 2013;16(12):1115-20.
  45. Qian LQ, Wang B, Niu JY, Gao S, Zhao DY. Assessment of the clinical effect of Chinese medicine therapy combined with psychological intervention for treatment of patients of peri-menopausal syndrome complicated with hyperlipidemia. *Chinese journal of integrative medicine*. 2010;16(2):124-30.
  46. Su JY, Xie QF, Liu WJ, Lai P, Liu DD, Tang LH, et al. Perimenopause Amelioration of a TCM Recipe Composed of *Radix Astragali*, *Radix Angelicae Sinensis*, and *Folium Epimedii*: An In Vivo Study on Natural



- Aging Rat Model. *Evid-Based Compl Alt*. 2013.
47. Wang LH, Wang X, Yu XZ, Xu WT. Potent therapeutic effects of shouwu jiangqi decoction on polycystic ovary syndrome with insulin resistance in rats. *Chinese journal of integrative medicine*. 2016;22(2):116-23.
  48. He Y, Li Q, Su M, Huang W, Zhu B. Acetylshikonin from Zicao exerts antifertility effects at high dose in rats by suppressing the secretion of GTH. *Biochemical and biophysical research communications*. 2016;476(4):560-5.
  49. Wei Y, Wang X, Yu Z, Zhou W, Wang L, Qin F, et al. Efficacy and Mechanism of Action of Yiru Tiao Jing Granule Against Hyperprolactinemia In Vitro and In Vivo. *Planta medica*. 2015;81(14):1255-62.
  50. Wu GL, Wu NY, Li TY, Fan YS, Yu GY, Lu WW. Effect of Chinese herbal medicines for nourishing yin, supplementing qi, and activating blood on reproductive endocrine activity and immune functions in patients with primary Sjogren's syndrome. *Chinese journal of integrative medicine*. 2015;21(10):778-83.
  51. Xia T, Fu Y, Li S, Ma R, Zhao Z, Wang B, et al. Bu Shen Tiao Chong recipe restores diminished ovary reserve through the BDNF pathway. *Journal of assisted reproduction and genetics*. 2016;33(6):795-805.
  52. Xie JH, Chen ZW, Pan YW, Luo DM, Su ZR, Chen HM, et al. Evaluation of safety of modified-Danggui Buxue Tang in rodents: immunological, toxicity and hormonal aspects. *Journal of ethnopharmacology*. 2016;183:59-70.
  53. Xu Y, Ma XP, An JN, Zhang ZJ, Ding J, Qu YK, et al. Short-time Qi Bao Mei Ran Formula Treatment Exerts Estrogenic Activities without Side Effects on Reproductive Tissues in Immature Mice. *Scientific reports*. 2015;5:17436.
  54. Zeng GL, Han XH, Yu J, Wang YH, Tian ZZ. Effect of Nourishing "Yin" Removing "Fire" Chinese Herbal Mixture on Hypothalamic Mammalian Target of Rapamycin Expression during Onset of Puberty in Female Rats. *Evid-Based Compl Alt*. 2015.
  55. Xu Y, Zhang ZJ, Geng F, Su SB, White KN, Blich SWA, et al. Treatment with Qing'E, A Kidney-Invigorating Chinese Herbal Formula, Antagonizes the Estrogen Decline in Ovariectomized Mice. *Rejuvenation research*. 2010;13(4):479-88.
  56. Zhou J, Liang Y, Lin XM, Ma RJ, Fang JQ. Is the efficacy of Chinese herbal medicine on polycystic ovarian syndrome enhanced by acupuncture in rats. *Biomed Res-India*. 2012;23(3):329-32.
  57. Mahesh VB. Hirsutism, virilism, polycystic ovarian disease, and the steroid-gonadotropin-feedback system: a career retrospective. *American journal of physiology Endocrinology and metabolism*. 2012;302(1):E4-E18.
  58. Reed BG, Carr BR. The Normal Menstrual Cycle and the Control of Ovulation. In: De Groot LJ, Chrousos G, Dungan K, Feingold KR, Grossman A, Hershman JM, et al., editors. *Endotext*. South Dartmouth MA: MDTText.com, Inc.; 2000.
  59. Burger HG, Dudley EC, Hopper JL, Groome N, Guthrie JR, Green A, et al. Prospectively measured levels of serum follicle-stimulating hormone, estradiol, and the dimeric inhibins during the menopausal transition in a population-based cohort of women. *The Journal of clinical endocrinology and metabolism*. 1999;84(11):4025-30.
  60. Randolph JF, Jr., Sowers M, Bondarenko IV, Harlow SD, Luborsky JL, Little RJ. Change in estradiol and follicle-stimulating hormone across the early menopausal transition: effects of ethnicity and age. *The Journal of clinical endocrinology and metabolism*. 2004;89(4):1555-61.
  61. Badawy A, Wageah A, El Gharib M, Osman EE. Prediction and Diagnosis of Poor Ovarian Response: The Dilemma. *Journal of Reproduction & Infertility*. 2011;12(4):241-8.
  62. Fan JX, Zhang B, Li LL, Xiao CW, Oladele OA, Jiang GL, et al. Effect of Soyabean Isoflavones Exposure on Onset of Puberty, Serum Hormone Concentration and Gene Expression in Hypothalamus, Pituitary Gland and Ovary of Female Bama Miniature Pigs. *Asian Austral J Anim*. 2015;28(11):1573-82.
  63. Wang DF, Zhang NY, Peng YZ, Qi DS. Interaction of zearalenone and soybean isoflavone on the development of reproductive organs, reproductive hormones and estrogen receptor expression in prepubertal gilts. *Animal reproduction science*. 2010;122(3-4):317-23.
  64. Abdi F, Mobedi H, Roozbeh N. Hops for Menopausal Vasomotor Symptoms: Mechanisms of Action. *Journal of menopausal medicine*. 2016;22(2):62-4.



65. Azizi H, Feng Liu Y, Du L, Hua Wang C, Bahrami-Taghanaki H, Ollah Esmaily H, et al. Menopause-related symptoms: traditional Chinese medicine vs hormone therapy. *Alternative therapies in health and medicine*. 2011;17(4):48-53.
66. Borrione P, Rizzo M, Quaranta F, Ciminelli E, Fagnani F, Parisi A, et al. Consumption and biochemical impact of commercially available plant-derived nutritional supplements. An observational pilot-study on recreational athletes. *Journal of the International Society of Sports Nutrition*. 2012;9(1):28.